

11. Cronnolly L. A. Anaesthetic management of obstructive sleep apnea patients. *J Clin Anesth* 1991; 3: 461-9.
12. Seet E., Chung F. Management of sleep apnea in adults-functional algorithms for the perioperative period: continuing professional development. *Can J Anesth.* 2010; 57: 849-65.
13. Ostermeier A. M., Roizen M. F., Hautkappe M., Klock P. A., Klapka J. M. Three sudden post-operative respiratory arrests associated with epidural opioids in patients with sleep apnea. *Anesth Analg* 1997; 85: 452-60.
14. Gupta R.M., Parvizi J., Hanssen A.D. Postoperative complications in patients with obstructive sleep apnea syndrome undergoing hip or knee replacement: a case-control study. *Mayo Clin Proc.* 2001; 76: 897-905.
15. Liao P., Yegneswaran B., Vairavanathan S. Postoperative complications in patients with obstructive sleep apnea: a retrospective matched cohort study. *Can J Anaesth.* 2009; 56: 819-28.
16. Squadrone V., Cocha M., Cerutti E. Continuous positive airway pressure for treatment of post-operative hypoxemia: a randomized controlled trial. *JAMA* 2005; 293: 589-95.
17. Ferreyra G.P., Baussano I., Squadrone V. Continuous positive airway pressure for treatment of respiratory complications after abdominal surgery: a systematic review and metaanalysis. *Ann Surg.* 2008; 247: 617-26.

Надійшла 7.10.2016

Рецензент д-р мед. наук, проф. О. О. Тарабрін

UDC 615.212.7.03:617-089.168.1

B. Tuyakov, Z. Stepien, D. Onichimowski, E. Mayzner-Zawadzka

## THORACIC WALL NERVE BLOCK

*Department Anesthesiology and Intensive Care, Clinical Hospital WSS,  
Olsztyn, Poland,*

*Department Anesthesiology and Intensive Care, Medical Faculty,  
University of Warmia and Mazury, Olsztyn, Poland*

УДК 615.212.7.03:617-089.168.1

**Б. Туяков, З. Степьян, Д. Оничимовский, Е. Майзнер-Завадска  
БЛОКАДЫ НЕРВОВ ГРУДНОЙ СТЕНКИ**

Блокады грудной клетки являются простым и эффективным методом лечения боли, характеризуются низким риском развития побочных эффектов и осложнений. Они требуют несколько точек размещения иглы и частой ориентации иглы для просмотра необходимой локализации. При этих типах блокад местный анестетик достигает нервов из интерфасциального пространства. Благодаря поверхностному расположению данных пространств мы можем использовать высокочастотные ультразвуковые датчики с высоким разрешением. Число точек приложения местных анестетиков постоянно увеличивается, что является шагом к открытию нового поколения блокад основанных на ультразвуковом контроле. Предстоящие проспективное рандомизированное и контролируемые клинические испытания могут клинически доказать их эффективность в торакальной хирургии.

Блокады грудной клетки: PECS I, PECS II и межлестничная блокада выполняются с помощью ультразвукового исследования при операциях на груди, множественных переломах ребер и других процедурах на грудной клетке. В данной работе описаны методы и блок анатомии. Кроме того, обсуждаются показания, осложнения и будущие перспективы блоков.

**Ключевые слова:** торакальная хирургия, региональная анестезия, PECS I, PECS II блокада, послеоперационное лечение болевого синдрома.

UDC 615.212.7.03:617-089.168.1

**B. Tuyakov, Z. Stepien, D. Onichimowski, E. Mayzner-Zawadzka  
THORACIC WALL NERVE BLOCK**

Due to the widespread access to ultrasonography, regional anaesthesia has become a safe method of post-operative pain management. It can be performed not only in orthopedic surgeries, but also in thoracic wall surgeries. One of the most frequent chest wall operations are breast surgeries. Although thoracic epidural anaesthesia and paravertebral blockades continue to be relevant, increasing number of one-day procedures demands an easier alternative with a lower number of side effects. To meet these expectations, in 2011 R. Blanco introduced pectoral nerve blocks. In PECS I block local anaesthetic is injected between the pectoral major and minor muscles. PECS II block extends analgesia to the axilla and consists of PECS I block and a second injection of local anaesthetic between the pectoral minor and serratus anterior muscles. Another effective method of antero-lateral chest analgesia is serratus plane block, where LA is deposited in the interfascial plane above the serratus muscle. Above-mentioned procedures lead to a lower opioid consumption after chest surgery and, as a consequence, a shorter stay in PACU.

**Key words:** Breast surgery; Regional anaesthesia; PECS I block; PECS II block; Serratus plane block; Post-operative pain management.

Regional anaesthesia allows effective blocks to be performed not only to the limbs, but also to the thoracic wall nerves. The progress that has been made in this discipline over the last decade has been possible thanks to the widespread availability of ultrasonography, which has led to a renaissance in regional anaesthesia. Regional techniques have evolved to make anaesthesia increasingly effective. The history of anaesthesia is full of cases when the modification of one small detail of a technique — an anatomical point or the angle of placing the needle — allowed the author to give his name to the approach. The pectoral block has a different genesis.

In 2011 Rafael Blanco proposed a block that has no analogy with the other approaches (R. Blanco, 2011). It is only similar to the transversus abdominis plane block (TAP block), where the local anaesthetic is placed into the interfascial plane between an internal oblique muscle and a transverse abdominal muscle (A. Kumar, 2015). In an article published in the 2011 edition of *Anesthesia* the author described an easy alternative to the thoracic epidural anaesthesia and thoracic paravertebral blockade in the post-operative pain management after breast surgery.

Breast cancer is the most common cancer among women. In the United States it affects one in eight of the female population. Effective pain control is a major challenge (R. Blanco et al., 2013; B. Fox, 2015). An increasing number of surgeries are being performed as a day procedures, making thoracic epidural anaesthesia or paravertebral blockade inadequate for these operations. Moreover, the side effects of these blocks could exceed the advantages. To solve this problem, the PECS blocks were introduced (R. Blanco et al., 2012; R. Sedra, 2015).

A PECS block derives its name from the word “pectoral” — the local anaesthetic is deposited in the fascial plane between the pectoral minor and major muscles. In Blanco’s study, the blockade was performed in approximately 50 patients, who required only minimal post-operative analgesia — a regular administration of paracetamol and dexketoprofen (R. Blanco, 2011). The block is useful for reconstructive breast cancer surgery or subpectoral prosthesis. It is performed with a linear ultrasound probe placed in a position similar to that used to the infraclavicular plexus block. The needle is inserted on the cephalad side of the probe just beneath the clavicle (M. Saleem, V. Irvine). First, the pectoral major muscle is identified. Then, between both pectoral muscles, the pectoral branch of thoraco-acromial artery is localised with a colour Doppler. The lateral pectoral nerve is located adjacent to the thoraco-acromial artery. The medial pectoral nerve is situated in the same interfascial space. In Blanco’s study 50 ml of the local anaesthetic (LA) 0.25 % bupivacaine — was injected. In subsequent publications, the LA amount was lim-

ited to 10–20 ml. The location was suitable to leave a catheter and perform a continuous analgesia using 0,25 % bupivacaine 5ml/h during for 7 days. Most of the patients with continuous analgesia did not require any opioids.

In 2012 Blanco described a modification to the PECS bloc, which was named the PECS II block (R. Blanco et al., 2012). He analysed the anatomy of the breast innervation, which was essential for the use of ultrasonography. This modification was intended to extend analgesia to the axilla, which provides a better pain control after vast revisions, tumorectomies, mastectomies and sentinel node dissection (Abrahams et al., 2016; R. Sedra, 2015). To perform a PECS II block two needle approaches are needed. The first approach is a PECS I blockade with the insertion of 10 ml of the LA. The second puncture injects 20 ml of local anaesthetic between pectoral minor muscle and serratus muscle. This enables reaching so called “axillary door” and blocks the long thoracic nerve and I and II intercostal nerves (R. Blanco et al. ; 2012). This approach also blocks the lateral branches of the intercostal nerves, which exit at the level of the mid-axillary line to innervate the mammary gland and the skin from T2 to T6.

### **Anatomy**

The pectoral muscles are mainly innervated by the lateral and medial pectoral nerves, which arise from the brachial plexus (R. Blanco et al., 2012). The lateral pectoral nerve arises from C5, C6 and C7 and is separated from the lateral cord of brachial plexus. It runs between the major and minor pectoral muscles in close proximity to the pectoral branch to the thoracoacromial artery. It innervates two-thirds of the pectoral major muscle. Medial pectoral nerve arises from C8-T1 and is separated from the medial cord of brachial plexus, running under pectoral minor muscle piercing it and the clavipectoral fascia and innervates the lower third of pectoral major muscle. It is possible that the medial pectoral nerve runs along the lateral border of the pectoral minor muscle.

A second group of nerves are the thoracic intercostal nerves from T2 to T6 (H. Ueshima et al., 2016), which run in a plane between the intercostal nerves to the sternum. The anterior divisions of these nerves pierce through the intercostal muscles, intercostal membrane and pectoral major muscle crossing the internal mammary artery and supply the medial side of breast. Lateral branches of intercostal nerves pierce intercostal muscles and the serratus anterior muscle in the mid-axillary line, giving off anterior and posterior terminal branches. The lateral branch of the intercostal nerve T2 continues as an intercostobrachial nerve to the axillary area.

A third group of nerves are the long thoracic and thoracodorsal nerves. The long thoracic nerve arises from C5-C7, enters the axilla runs along the serratus anterior muscle. Once damaged during the radical mastectomy or axillary clearance, a winging scapula can be produced, especially when the arm is lifted forward. Damage to the long thoracic nerve has also been described during the plexus brachialis blockade from the interscalene approach, when the needle was inserted through the middle scalene muscle.

The thoracodorsal nerve is a branch of the posterior trunk of the brachial plexus. It runs along the thoracodorsal artery and innervates the latissimus dorsi muscle (the posterior axilla wall). The nerve lies very deep and can be damaged during the breast reconstruction using serratus anterior flaps.

The clavipectoral fascia lies on the anterior surface of pectoral minor nerve and on the lateral border of the muscle converts to Gerdy ligament (the suspensory ligament of axilla). It enables maintaining the concave shape of the axilla.

### **Sonoanatomy and block technique**

#### *PECS I block*

The patient is positioned supine with the arm put on the side or abducted and externally rotated. Blanco described an approach when the probe is located at the level of the


coracoid process. Perez has modified the probe placement to the lateral one-third of the clavicle (M. F. Perez et al., 2013). The needle is inserted medial-to-lateral in the “in-plane” position to minimize the possibility of the bone or vascular structure damage. The following structures must be identified: subcutaneous tissue, pectoral major muscle, pectoral minor muscle, axillary artery, axillary vein and pleura (E. D. Bolin et al., 2015). Between both pectoral muscles there is the thoracoacromial artery and the lateral pectoral nerve. 10–20 ml of 0.125–0.25 % bupivacaine or ropivacaine is injected. This block does not affect the anterior branches of the intercostal nerves (Fig. 1).

#### *PECS II block*

The initial part is similar to a PEC 1 block (Abhijit S. Nair et al., 2015). Then the probe is moved laterally and distally until the third rib (J. S. Kim et al., 2016) and the lateral border of the pectoral minor muscle are identified. On the third rib, continuation of the Gerdy ligament is visualized along with serratus anterior muscle, which lays beneath the ligament and covers the ribs (E. D. Bolin et al., 2015). The parietal pleura is located at the depth of serratus muscle between the ribs. The needle is inserted “in-plane”, medially to laterally and 20 ml of long lasting local anaesthetic is administrated. During the withdrawal of the needle additional 10 ml of local anaesthetic can be injected into the interfascial plane between two pectoral muscles. Thanks to this method, the long thoracic nerve, thoracodorsal nerve and intercostal nerves II–IV are blocked. In the PECS II block the LA spreads along the chest wall until the level of T8 (Fig. 2).

#### *Serratus Plane Block*

A serratus plane block provides an exact analgesia of the thoracic wall by blocking the lateral branches of intercostal nerves (M. P. Sebastian 2014; H. Otake, 2015; J. S. Kim et al., 2016,) . The blockade is performed in the supine position. We visualize the V



*Fig. 1. PECS I block. The plane between pectoral muscles is visible. The tip of needle is so deep into pectoral minor muscle and should be remove some milimetres. LA injection should be given exactly into plane between pectoral muscles. Arrow 1 — fascial plane between pectoral muscles*


rib at the mid-axillary line, latissimus dorsi muscle (more superficially and laterally), teres major muscle (above) and serratus anterior muscle. The needle should be inserted to a depth of about 2 cm . The local anaesthetic is injected into the interfascial plane above the serratus muscle or beneath this muscle. According to Blanco placing the LA above the muscle provides a faster sensory blockade of the hemi thorax(R. Blanco et al., 2013). The thoracodorsal artery can serve as a landmark to identify the localization. Superficial placement of LA is one of the advantages of this technique(P. A.-de la Torre et al., 2015; S. Tighe et al., 2013) (Fig. 3).

### Practical Use


A PECS I block can be an effective method for postoperative analgesia for mastectomies, breast augmentation, breast reconstruction, CRTD implantation (Fujiwara et al., 2014), or vascular port implantation. PECS II indications are similar to a PEC I's plus mastectomy with axillary clearance, sentinel node dissection and anterior thoracotomy.

A serratus-plane block is useful as an additional analgesia in breast surgery, multiple rib fractures (N. P. Kunhabdulla et al., 2014) (Fig. 4) and breast reconstructive surgeries involving the latissimus dorsi muscle (Abhijit S. Nair et al., 2015). It is also used as anterolateral chest chronic pain treatment (Fujiwara S. et al., 2015) after radiotherapy and minimal invasive cardiac surgeries (MIDCAB — minimal invasive direct coronary artery by-pass) .

Initially, few cases of opioid reduction in these blocks were published. In 2015 G. M. Bashandy (G. M. Bashandy et al., 2015) with colleagues published a PEC II prospective randomized trial, in which 120 patients underwent a modified unilateral radical mastectomy. They were divided into two groups — with and without the block. In patients with a PECS II block the opioid consumption was reduced during the first 12 hours after sur-



*Fig. 2.* PECS II block. The end-point of the needle is between pectoral minor muscle and serratus anterior muscle. Arrow 1 — fascial plane between two muscles. Arrow 2 — the shift of the needle



*Fig. 3.* Sonoanatomy by Serratus Plane Block. The probe is placed in the middle axillary line in the short axis. Arrows *I* — the plane between two muscles

gery, there was a 50 % reduction of fentanyl use during the operation, a shorter stay in a post-anaesthesia care unit and a generally shorter stay in the hospital.

S. S. Wahba compared thoracic paravertebral block (PVB) versus a PECS II block in the radical mastectomies (S. S. Wahba et al., 2013). The PECS II block required lower opioid consumption during the first 24 hours after the surgery. After the PECS II block also pain intensity was also lower during the first 12 hours compared to PVB. The PONV was comparable between the two groups. The authors suggest that a PECS II block can be an alternative to a paravertebral block. It is a safe, effective chest wall block method with fewer contraindications connected with antithrombotic therapy compared to epidural thoracic analgesia and, unlike PVB, PECS II also provides a contralateral sympathetic chain block.

In 2016, Abrahams published a review of several truncal block methods (M. Abrahams et al, 2016). It was noted that current publications recommend thoracic cage blocks with an “A” grade of recommendation and Ib-III level of evidence.


All of the cited studies described these blockades as an intra- and postoperative analgesia, which was accompanied by general anaesthesia (A. Amir et al., 2016). There are only few examples of using them as the only one method of anaesthesia (H. Murata et al., 2015).

### Complications

Few complications of the block have been described (A. J. Louw, 2014). The thoraco-acromial artery should be localized to avoid a vascular LA administration. In the PECS II block the needle must be visualized to avoid pleura puncture. The chest wall blockades are performed with a large volume of local anaesthetic, so the maximal doses should be followed, taking into account the age and a general condition of the patient to prevent any toxic reactions.

### Summary

Chest wall blockades are an easy and effective method of pain management, characterized by a low risk of side effects and complications. They require a several points of needle placement and a frequent needle orientation to view the desired localization. In these types of blocks, the local anaesthetic reaches nerves from the interfascial plane. Thanks to the superficial site of these spaces, high frequency ultrasound probes can be used, which provides a better image resolution. An increasing number of new compartments of local anaesthetic placement have been introduced, which is a step toward pre-



*Fig. 4.* Continuous Serratus Plane Block with Certa Catheter (Ferrosand TM, Denmark) by polytrauma. The block end-point of the needle is performed for the patient with ribs multiple fractures (II–XI) . Additionally, the patient had spine injury with the fractures of left transverse processis of vertebrae Th II–Th10 and multiple fractures of pelvis bones . Thoracic paravertebral block or thoracic epidural analgesia were contraindicated due to thoracic vertebrae fractures. Lumbar spine was intacted. Lumbar epidural analgesia was performed for treatment of pelvic pain

senting a new generation of blockades based on ultrasound guidance. Upcoming, prospective, randomized and controlled trials may show a clinical evidence of their efficacy in breast and chest wall surgeries.

Chest wall blockade: PECS I, PECS II and serratus-plane block are performed using ultrasonography in breast perations, multiple rib fractures and other chest wall procedures. In this paper, block techniques and the anatomy and sonoanatomy of the breast area are described. Moreover, the indications, complications and future perspectives of the blocks are discussed.

#### ЛИТЕРАТУРА

1. *Bashandy G. M.* Pectoral nerves I and II blocks in multimodal analgesia for breast cancer surgery: a randomized clinical trial / G. M. Bashandy, D. N. Abbas // *Reg Anesth Pain Med.* – 2015. – Vol. 40. – P. 68–74.
2. *Blanco R.* The ‘pecs block’: a novel technique for providing analgesia after breast surgery / R. Blanco // *Anaesthesia.* – 2011. – Vol. 66. – P. 847–848.
3. *Blanco R.* Ultrasound description of Pecs II (modified PecsI): a novel approach to breast surgery / R. Blanco, M. Fajardo, T. P. Maldonado // *Rev Esp Anesthesiol Reanim.* – 2012. – Vol. 59. – P. 470–475.
4. *Blanco R.* Reply to the article entitled “Ultrasound description of Pecs II (modified Pecs I): a novel approach to breast surgery”. Reply of the authors / R. Blanco, T. P. Maldonado // *Rev Esp Anesthesiol Reanim.* – 2013. – Vol. 60. – P. 296–297.
5. *Serratus-intercostal plane block.* An encouraging approach for breast surgery / M. F. Perez, P. A. de la Torre, S. Lopez, F. J. Garcia // *Anaesthesia.* – 2012. – Vol. 66. – P. 1463.

6. *Perez M. F.* A new approach to pectoralis block / M. F. Perez, J. G. Miguel, P. A. de la Torre // *Anaesthesia*. – 2013. – Vol. 68. – P. 430.
7. *Sebastian M. P.* Pecs II or serratus plane block / M. P. Sebastian // *Anaesthesia*. – 2014. – Vol. 69. – P. 1173.
8. *Bolin E. D.* Regional Anesthesia for Breast Surgery: Techniques and Benefits / E. D. Bolin // *Curr. Anesthesiol Rep* June. – 2015. – Vol. 5. – Issue 2. – P. 217–224.
9. *Louw A. J.* Procedural sedation and analgesia for out-of-hospital breast surgery: an overview of the procedural sedation and analgesia technique / A. J. Louw // *Southern African Journal of Anaesthesia and Analgesia*. – 2014. – Vol. 20. – P. 89–92.
10. *De la Torre P. A.* Thoracic Paravertebral Block and Its Effects on Chronic Pain and Health-Related Quality of Life After Modified Radical Mastectomy / P. A. de la Torre, M. F. Perez // *Regional Anesthesia and Pain Medicine*. – 2015. – Vol. 40. – Issue 2. – P. 177–178.
11. *Tighe S. K.* Serratus plane block: do we need to learn another technique for thoracic wall blockade / S. K. Tighe, M. K. Karmakar // *Anaesthesia*. – 2013. – Vol. 68. – P. 1095–1106.
12. *Abrahams M.* Update on Ultrasound for Truncal Blocks. A Review of the Evidence / M. Abrahams, R. Derby, J.-L. Horn // *Reg Anesth Pain Med*. – 2016. – Vol. 41. – P. 275–288.
13. *Wahba S. S.* Thoracic paravertebral block versus pectoral nerve block for analgesia after breast surgery / S. S. Wahba, S. M. Kamal // *Egyptian J Anaesth*. – 2014. – Vol. 30. – P. 129–135.
14. *Kunhabdulla N. P.* Serratus Anterior Plane Block for Multiple Rib Fractures / N. P. Kunhabdulla, A. Agarwal // *Pain Physician*. – 2014. – Vol. 17. – P. 65–662.
15. *Ueshima H.* Serratus Plane Block for a Contraction of the Latissimus Dorsi Muscle / H. Ueshima, A. Kitamura // *Regional Anesthesia and Pain Medicine*. – 2016. – Vol. 41, N 3. – P. 411.
16. *Implementation of Pecs I and Pecs II Blocks as Part of Opioid-Sparing Approach to Breast Surgery / A. Amir, S. Jolin, S. Amberg, S. Nordstrom // Regional Anesthesia and Pain Medicine. – 2016. – Vol. 41, N 4. – P. 544–545.*
17. *Sedra R.* Pectoralis Minor Nerve Block versus Thoracic Epidural and Paravertebral Block in Perioperative Pain Control of Breast Surgery (Mini Review) / R. Sedra // *Journal of Anesthesiology and Critical Care Medicine*. – 2015. – Vol. 2. – P. 1–4.
18. *Ultrasound guided blocks for surgeries: procedures involving chest wall (Pecs 1, 2 and serratus plane block) / A. S. Nair, R. K. Sahoo, M. Ganapathy, R. Mudunuri // Anesthesia, Pain & Intensive Care. – 2015. – Vol. 19, N 3. – P. 348–351.*
19. *Murata H.* Pecs block for anesthesia in breast surgery of the elderly / H. Murata, T. Ichinomiya, T. J. Hara // *Anaesthesia*. – 2015. – Vol. 29. – P. 644.
20. *Plane Block and Ultrasound-Guided Intermediate Cervical Plexus Block After Single-Incision Transaxillary Robotic Thyroidectomy. A Prospective, Randomized, Controlled Trial / S. Kim, E.-Y. Soh, H. Ahn [et al.] // Regional Anesthesia and Pain. – 2016. – Vol. 41, N 5. – P. 21–38.*
21. *Fujiwara A.* Pectoral nerves (PECS) and intercostal nerve block for cardiac resynchronization therapy device implantation / A. Fujiwara, N. Komasa, T. Minami // *Springerplus*. – 2014. – Vol. 3. – P. 409.
22. *Serratus plane block: a novel ultrasound-guided thoracic wall nerve block // R. Blanco, T. Parras, J. G. McDonnell, A. Prats-Galino // Anaesthesia. – 2013. – Vol. 68. – P. 1107–1113.*
23. *Fujiwara S.* Pectoral nerve blocks and serratus–intercostal plane block for intractable postthoracotomy syndrome / S. Fujiwara, N. Komasa, T. Minami // *Journal Clin Anesth*. – 2015. – Vol. 27. – P. 275–276.
24. *Saleem M.* Pain Score “0/10” A novel approach for free pain breast surgery : Highlights and Abstracts XXXIV ESRA Congress, Lubljana, Sept. 2014 / M. Saleem, V. Irvine. – 2014. – P. 1–16.
25. *Ultrasound guided blocks for surgeries: procedures involving chest wall (Pecs 1, 2 and serratus plane block) / A. S. Nair, R. K. Sahoo, M. Ganapathy, R. Mudunuri // Anaesth Pain & Intensive Care. – 2015. – Vol. 19, N 3. – P. 348–351.*

## REFERENCES

1. Bashandy G.M., Abbas D.N. Pectoral nerves I and II blocks in multimodal analgesia for breast cancer surgery: a randomized clinical trial. *Reg Anesth Pain Med*. 2015; 40: 68-74.
2. Blanco R. The ‘pecs block’: a novel technique for providing analgesia after breast surgery. *Anaesthesia* 2011; 66: 847-848.
3. Blanco R., Fajardo M., Parras Maldonado T. Ultrasound description of Pecs II (modified PecsI): a novel approach to breast surgery. *Rev Esp Anesthesiol Reanim*. 2012; 59: 470-475.



4. Blanco R., Parras Maldonado T. Reply to the article entitled "Ultrasound description of Pecs II (modified Pecs I): a novel approach to breast surgery". Reply of the authors. *Rev Esp Anesthesiol Reanim.* 2013; 60: 296-297.
5. Perez Fajardo M., Alfaro de la Torre P., Lopez S., Garcia F.J. Serratus-intercostal plane block. An encouraging approach for breast surgery. *Anaesthesia* 2012; 66: 1463.
6. Perez MF., Miguel JG., de la Torre PA. A new approach to pectoralis block. *Anaesthesia.* 2013; 68: 430.
7. Sebastian M.P. Pecs II or serratus plane block. *Anaesthesia* 2014; 69: 1173.
8. Eric D. Bolin Regional Anesthesia for Breast Surgery: Techniques and Benefits. *Curr. Anesthesiol Rep* June 2015; 5 (2): 217-224
9. Louw AJ. Procedural sedation and analgesia for out-of-hospital breast surgery: an overview of the procedural sedation and analgesia technique. *Southern African Journal of Anaesthesia and Analgesia* 2014; 20: 89-92
10. de la Torre P.A., Perez A. Thoracic Paravertebral Block and Its Effects on Chronic Pain and Health-Related Quality of Life After Modified Radical Mastectomy. *Regional Anesthesia and Pain Medicine.* March-April 2015; 40 (2): 177-178
11. Tighe S.K., Karmakar M.K. Serratus plane block: do we need to learn another technique for thoracic wall blockade. *Anaesthesia* 2013; 68: 1095-1106.
12. Abrahams M., Derby R., Horn J.-L. Update on Ultrasound for Truncal Blocks. A Review of the Evidence. *Reg Anesth Pain Med* 2016; 41: 275-288.
13. Wahba S.S., Kamal S.M. Thoracic paravertebral block versus pectoral nerve block for analgesia after breast surgery. *Egyptian J Anaesth.* 2014; 30: 129-135.
14. Kunhabdulla N.P., Agarwal A. Serratus Anterior Plane Block for Multiple Rib Fractures Pain. *Physician* 2014; 17: 65-662.
15. Ueshima H., Kitamura A. Serratus Plane Block for a Contraction of the Latissimus Dorsi Muscle. *Regional Anesthesia and Pain Medicine* May-June 2016; 41; 3: 411.
16. Amir A., Jolin S., Amberg S., Nordstrom S. Implementation of Pecs I and Pecs II Blocks as Part of Opioid-Sparing Approach to Breast Surgery. *Regional Anesthesia and Pain Medicine.* July-August 2016; 41; 4: 544-545.
17. Sedra R. Pectoralis Minor Nerve Block versus Thoracic Epidural and Paravertebral Block in Perioperative Pain Control of Breast Surgery — Mini Review. *Journal of Anesthesiology and Critical Care Medicine* 2015; 2: 1-4.
18. Nair A.S., Sahoo R.K., Ganapathy M., Mudunuri R. Ultrasound guided blocks for surgeries: procedures involving chest wall (Pecs 1, 2 and serratus plane block). *Anaesth Pain & Intensive Care* 2015; 19(3): 348-351.
19. Murata H., Ichinomiya T., Hara T. Pecs block for anesthesia in breast surgery of the elderly. *J Anesth.* 2015; 29: 644.
20. Jin-Soo Kim, Euy-Young Soh, Hyoeun Ahn, Sang Eon Oh, Jung-Dong Lee, Han Bum Joe. Plane Block and Ultrasound-Guided Intermediate Cervical Plexus Block After Single-Incision Transaxillary Robotic Thyroidectomy. A Prospective, Randomized, Controlled Trial. *Regional Anesthesia and Pain Medicine.* September-October 2016; 41 (5): 21-38.
21. Fujiwara A., Komazawa N., Minami T. Pectoral nerves (PECS) and intercostal nerve block for cardiac resynchronization therapy device implantation. *Springerplus* 2014; 3: 409.
22. Blanco R., Parras T., McDonnell J.G., Prats-Galino A. Serratus plane block: a novel ultrasound-guided thoracic wall nerve block. *Anaesthesia* 2013; 68: 1107-1113.
23. Fujiwara S., Komazawa N., Minami T. Pectoral nerve blocks and serratus-intercostal plane block for intractable postthoracotomy syndrome. *Journal Clin Anesth.* 2015; 27: 275-276.
24. Saleem M., Irvine V. Pain Score "0/10" A novel approach for free pain breast surgery. Highlights and Abstracts XXXIV ESRA Congress, Lubljana, Sept. 2014: 1-16.
25. Nair AS., Sahoo R.K., Ganapathy M., Mudunuri R. Ultrasound guided blocks for surgeries: procedures involving chest wall (Pecs 1, 2 and serratus plane block). *Anaesth Pain & Intensive Care* 2015; 19(3): 348-351.

Надійшла 7.10.2016

Рецензент д-р мед. наук, проф. О. О. Тарабрін